

DSM-IV DIAGNOSTIC CHECKLIST

Rater: _____

Circle one:

1. Examiner 2. Diagnostician

Consensus Document:

1. Yes 2. No

Reviewed at JHH

1. Yes 2. No

Date: ____/____/____

Date Revised: ____/____/____

SEPARATION ANXIETY DISORDER

	<u>No</u>	<u>Yes</u>	<u>DK</u>	
A.				
Developmentally inappropriate and excessive anxiety concerning separation from home or from those to whom the individual is attached, as evidenced by three (or more) of the following:				
(1)	1	3	9	(46)
(2)	1	3	9	(47)
(3)	1	3	9	(48)
(4)	1	3	9	(49)
(5)	1	3	9	(50)
(6)	1	3	9	(51)
(7)	1	3	9	(52)
(8)	1	3	9	(53)
Criterion A = 3, if A1 through A8 ≥ 3	1	3	9	(54)
If Criteria A = 1, Circle Separation Anxiety Disorder = 1, and Skip				
B.	1	3	9	(55)
C.	1	3	9	(56)
D.	1	3	9	(57)
E.	1	3	9	(58)

Circle Separation Anxiety Disorder = 3, if Criteria A through E = 3

SEPARATION ANXIETY DISORDER.....1.....2.....3.....9 (59)

Age of Onset, if 2 or 3: _____ (60-61)

ATTENTION-DEFICIT HYPERACTIVITY DISORDER

A.I. <u>INATTENTION</u>	<u>No</u>	<u>Yes</u>	<u>DK</u>
1. Difficulty Sustaining Attention on Tasks or Play	1	3	9
2. Easily Distracted	1	3	9
3. Makes a lot of Careless Mistakes	1	3	9
4. Doesn't Listen	1	3	9
5. Difficulty Following Instructions	1	3	9
6. Difficulty Organizing Tasks	1	3	9
7. Dislikes/Avoids Tasks Requiring Attention	1	3	9
8. Loses Things	1	3	9
9. Forgetful in Daily Activities	1	3	9

CRITERION A.I. = 3; if 1 through 9 ≥ 6 1 3 9

A.II. HYPERACTIVITY/IMPULSIVITY

1. Difficulty Remaining Seated	1	3	9
2. Impulsivity	1	3	9
3. Fidgets	1	3	9
4. Runs or Climbs Excessively	1	3	9
5. On the go/Acts as if Driven by a Motor	1	3	9
6. Difficulty Playing Quietly	1	3	9
7. Blurts Out Answers	1	3	9
8. Difficulty Waiting Turn	1	3	9
9. Often Interrupts or Intrudes	1	3	9
10. Talks Excessively	1	3	9

ATTENTION-DEFICIT HYPERACTIVITY DISORDER (CONTINUED)

	<u>No</u>	<u>Yes</u>	<u>DK</u>
CRITERION A.II = 3 IF 1 and 3 through 10 ≥ 6 (not 2)	1	3	9
A: AI present and AII present	1	3	9
B. Duration of Symptoms 6 months	1	3	9
C. Some symptoms that caused impairment present before the age of 7:	1	3	9
D. Some impairment from symptoms must be present in two or more situations (e.g. School and home)	1	3	9
E. Clinically significant impairment	1	3	9
F. Does not meet criteria for Pervasive Developmental Disorder	1	3	9

Circle ADHD = 3 if criteria A through F are met.

ATTENTION-DEFICIT HYPERACTIVITY DISORDER.....1.....2.....3.....9

Age of Onset, if 2 or 3: _____

Circle ADHD1 = 3 if subject meets Criterion A.I but not criterion A.II; and B through F = 3.

ADHD1: PREDOMINANTLY INATTENTIVE TYPE1.....2.....3.....9

Circle ADHD2 = 3 if subject meets criterion A.II but not criterion A.I; and B through F = 3.

ADHD2: PREDOMINANTLY HYPERACTIVE-IMPULSIVE TYPE1.....2.....3.....9

Circle ADHD3 = 3 if both criterion A.I and A.II are met; and B through F = 3.

ADHD3: COMBINED TYPE1.....2.....3.....9

COMPLETE ONLY FOR SUBJECTS AGED LESS THAN SIXTEEN

OPPOSITIONAL DEFIANT DISORDER

	<u>No</u>	<u>Yes</u>	<u>DK</u>
A. A pattern of negativistic, hostile and defiant behavior lasting at least 6 months, during which four (or more) of the following are present:	1	3	9
1) often loses temper	1	3	9
2) often argues with adults	1	3	9
3) often actively defies or refuses to comply with adults' requests or rules	1	3	9
4) often deliberately annoys people	1	3	9
5) often blames others for his or her mistakes or misbehavior	1	3	9
6) is often touchy or easily annoyed by others	1	3	9
7) is often angry and resentful	1	3	9
8) is often spiteful or vindictive	1	3	9
B. The disturbance in behavior causes clinically significant impairment in social, academic or occupational functioning	1	3	9
C. The behaviors do not occur exclusively during the course of Psychotic or Mood Disorder	1	3	9
D. Criteria are not met for Conduct Disorder and, if the individual is age 18 years or older, criteria are not met for Antisocial Personality Disorder.	1	3	9

Circle Oppositional Defiant Disorder = 3 if criteria A through D are met.

OPPOSITIONAL DEFIANT DISORDER.....1.....2.....3.....9

If Oppositional Defiant Disorder = 2 or 3, age at onset: _____

CONDUCT DISORDER

	<u>No</u>	<u>Yes</u>	<u>DK</u>
A. A repetitive and persistent pattern of behavior in which the basic rights of others or major age-appropriate societal norms or rules are violated as manifested by the presence of three (or more) of the following criteria in the past 12 months, with at least one criterion present in the past 6 months:	1	3	9
Aggression to People and Animals			
1) often bullies, threatens or intimidates others	1	3	9
2) often initiates physical fights	1	3	9
3) has used a weapon that can cause serious physical harm to others (e.g., a bat, brick, broken bottle, knife, gun)	1	3	9
4) has been physically cruel to people	1	3	9
5) has been physically cruel to animals	1	3	9
6) has stolen while confronting a victim (e.g., mugging, purse snatching, extortion, armed robbery)	1	3	9
7) has someone into sexual activity	1	3	9
Destruction of Property			
8) has deliberately engaged in fire setting with the intention of causing serious damage	1	3	9
9) has deliberately destroyed others' property (other than by fire setting)	1	3	9
Deceitfulness or Theft			
10) has broken into someone else's house, building or car	1	3	9
11) often lies to obtain goods or favors or to avoid obligations (i.e., "cons" others)	1	3	9
12) has stolen items of nontrivial value without confronting a victim (e.g., shoplifting, but without breaking and entering; forgery)	1	3	9
Serious Violations of Rules			
13) often stays out at night despite parental prohibitions, beginning before age 13 years	1	3	9
14) has run away from home overnight at least twice while living in parental or parental surrogate's home (or once without returning for a lengthy period)	1	3	9
15) is often truant from school, beginning before age 13 years	1	3	9
B. The disturbance in behavior causes clinically significant impairment in social, academic or occupational functioning	1	3	9
C. if the individual is age 18 years or older, criteria are not met for Antisocial Personality Disorder.	1	3	9

CONDUCT DISORDER (CONTINUED)

No Yes DK

Specify type based on age of onset:

Childhood-Onset Type: at least one criterion present prior to age 10 years 1 3 9

Adolescent-Onset Type: absence of any criterion prior to age 10 years 1 3 9

Circle Conduct Disorder = 3 if criteria A through C are met.

CONDUCT DISORDER.....1.....2.....3.....9

If Conduct Disorder = 2 or 3, age at onset: _____

MAJOR DEPRESSION

	<u>No</u>	<u>Yes</u>	<u>DK</u>	
A. <u>At least five</u> of the following symptoms have been present during the same two week period, with at least one being A1 or A2.	1	3	9	(62)

Elements of Criterion A:

1. Depressed or irritable mood	1	3	9	(64)
2. Diminished interest or pleasure	1	3	9	(65)
(If A1 and A2 both = 1, circle Criterion A = 1, Major Depression = 1 and skip to Mania)				
3. >5% Weight Change	1	3	9	(66)
4. Sleep Disturbance	1	3	9	(67)
5. Psychomotor Agitation or Retardation	1	3	9	(68)
6. Low Energy	1	3	9	(69)
7. Negative Self-Attitude	1	3	9	(70)
8. Impaired Concentration	1	3	9	(71)
9. Recurrent thoughts of death, or suicide attempt or plan	1	3	9	(72)
9a. Suicidal thoughts	1	3	9	
9b. Suicide plan	1	3	9	
9c. Suicide attempt	1	3	9	

CODE CRITERION A

	<u>No</u>	<u>Yes</u>	<u>DK</u>	
B. <u>NOT</u> coincident with a mixed episode.	1 coincident	3 not coincident	9	(73)
C. Significant distress or impairment	1	3	9	(81)
D. Not drug-induced or induced by a medical condition	1	3 not drug- or illness-induced	9	(82)

MAJOR DEPRESSION (CONTINUED)

	<u>No</u>	<u>Yes</u>	<u>DK</u>	
E. Not accounted for by bereavement (D.9)	1 accounted for	3 not accounted for	9	(83)
F. Not superimposed upon another psychosis (D.10)	1 superimposed	3 not superimposed	9	(84)

Circle Major Depression = 3 if all of A to F above are coded 3.

MAJOR DEPRESSION.....1.....2.....3.....9 (85)

If Major Depression = 2 or 3, age at first episode: ___ ___ (86-87)

If Major Depression = 2 or 3 number of episodes: ___ ___ ___ (88-90)

ADDITIONAL RATINGS FOR MAJOR DEPRESSION:

1. If Major Depression = 2, or 3,
specify the presence of psychotic features

1 absent	3 present	9	(91)
-------------	--------------	---	------

2. If psychotic features present, specify
mood congruency

1 mood congruent only	3 mood incongruent present	9	(92)
--------------------------------	-------------------------------------	---	------

MANIA

	<u>No</u>	<u>Yes</u>	<u>DK</u>	
A. At least one week of at least one of the following:	1	3	9	(99)
AA. At least four days of at least one of the following:	1	3	9	(100)

Elements of Criterion A:				
1. persistently elevated or expansive mood (E.1)	1	3	9	(102)
2. persistently irritable mood (E.1)	1	3	9	(103)
CODE CRITERION A				

(If Criterion A = 1, code Mania = 1 and skip to Dysthymic Disorder.)

<p>B. During the period of mood disturbance, at least three of the following symptoms have persisted (four if the mood is only irritable) and have been present to a significant degree</p>	1	3	9	(104)
---	---	---	---	-------

Elements of Criterion B:				
1. Increase in goal-directed activity (socially, at work or school, or sexually), or psycho-motor agitation	1	3	9	(105)
2. Hypertalkative or pressure to keep talking	1	3	9	(106)
3. Flight of ideas or subjective experience that thoughts are racing	1	3	9	(107)
4. Inflated self-esteem or grandiosity	1	3	9	(108)
5. Decreased need for sleep	1	3	9	(109)
6. Distractibility, i.e., attention too easily drawn to unimportant or irrelevant external stimuli	1	3	9	(110)
7. Excessive involvement in pleasurable activities which have a high potential for painful consequences, e.g., unrestrained buying sprees, sexual indiscretions, or foolish business investments	1	3	9	(111)
CODE CRITERION B				

MANIA (CONTINUED)

	<u>No</u>	<u>Yes</u>	<u>DK</u>	
C. <u>NOT</u> coincident with a mixed episode.	1	3 not coincident	9	(112)
D. One or both of the following is true:	1	3	9	(123)

Elements of Criterion D:

1. Symptoms cause marked impairment in occupational or social functioning	1	3	9	(124)
2. Psychotic features are present (in presence of mood change)	1	3	9	(125)

CODE CRITERION D

E. Not due to substance use or medical condition	1	3	9	(126)
	due to organic factor	not due to organic factor		

MANIA (CONTINUED)

	<u>No</u>	<u>Yes</u>	<u>DK</u>	
F. Mood symptoms are not better accounted for by Schizoaffective Disorder and are not superimposed on a psychotic disorder (i.e., no delusions or hallucinations in the absence of mood changes)	1 Superimposed	3 Not Superimposed	9	(127)
G. Unequivocal change in functioning	1	3	9	(128)
H. Change observable by others	1	3	9	(129)

Circle Mania = 3 if all criteria A through F are rated 3.

MANIA.....1.....2.....3.....9 (130)

If Mania = 2 or 3, age at onset: _____ (131-132)

If Mania = 2 or 3, number of episodes: _____ (133-135)

Circle Hypomania = 3, if AA, B, E, F, G, and H = 3 [D must = 1]

HYPOMANIA.....1.....2.....3.....9 (136)

If Hypomania = 2 or 3, age at onset: _____ (137-138)

If Hypomania = 2 or 3, number of episodes: _____ (139-141)

ADDITIONAL RATINGS FOR MANIA:

1. If psychotic features present, specify mood congruency	1 mood congruent only	3 mood incongruent present	9	(148)
---	--------------------------------	-------------------------------------	---	-------

MIXED EPISODE

	<u>No</u>	<u>Yes</u>	<u>DK</u>	
A. The criteria are met for both a Manic Episode and a Major Depressive Episode (except for duration) nearly every day during at least a one-week period	1	3	9	(149)

(If Criterion A = 1, code Mixed Episode = 1 and go directly to Dysthymic Disorder).

B. The mood disturbance is sufficiently severe to cause <u>at least one</u> of the following:	1	3	9	(150)
---	---	---	---	-------

Elements of Criterion B:				
1. marked impairment in occupational functioning or in usual social activities or relationships with others.	1	3	9	(151)
2. hospitalization to prevent harm to self or others.	1	3	9	(152)
3. psychotic features are present.	1	3	9	(153)
CODE CRITERION B.				

C. Not induced by substance or general medical condition.	1 induced	3 not induced	9	(154)
---	--------------	---------------------	---	-------

D. Not superimposed upon another psychosis.	1 superimposed	3 not superimposed	9	(155)
---	-------------------	--------------------------	---	-------

Code Mixed Episode = 3 of all of the above (Criteria A through D) are = 3.

MIXED EPISODE.....1.....2.....3.....9 (156)

If Mixed Episode = 2 or 3, code age at onset: _____ (157-158)

If Mixed Episode = 2 or 3, number of episodes: _____ (159-161)

DYSTHYMIC DISORDER

	<u>No</u>	<u>Yes</u>	<u>DK</u>	
A. Depressed mood more days than not, as indicated either by subjective account or observation by others, for at least 2 years.	1	3	9	(162)

If A = 1, then code Dysthymia = 1 and go to Psychosis

B. Presence while depressed of at least two of the following symptoms (1-6):	1	3	9	(163)
--	---	---	---	-------

Elements of Criterion B:				
1. Poor appetite or overeating	1	3	9	(164)
2. Insomnia or hypersomnia	1	3	9	(165)
3. Low energy or fatigue	1	3	9	(166)
4. Low self-esteem	1	3	9	(167)
5. Poor concentration or difficulty making decisions	1	3	9	(168)
6. Feelings of hopelessness	1	3	9	(169)
CODE CRITERION B				

C. Never symptom-free for two months	1	3	9	(175)
D. No Major Depressive episode during the first two years	1	3	9	(176)
E. No manic, mixed, or hypomanic episodes; not meet criteria for Cyclothymic Disorder	1	3	9	(177)
F. Not superimposed upon another psychosis	1	3	9	(178)
G. Not due to medical condition or substance use	1	3	9	(179)
H. Symptoms cause significant distress or impairment in functioning	1	3	9	(180)

Circle Dysthymic Disorder = 3 if all of criteria A through H are rated 3.

DYSTHYMIC DISORDER.....1.....2.....3.....9 (181)

If Dysthymic Disorder = 2 or 3, age at onset: _____ (182-183)

PSYCHOSIS/SCHIZOPHRENIA

A. Characteristic symptoms: Two (or more) of the following, each present for a significant portion of time during a 1-month period (or less if successfully treated):

	<u>No</u>	<u>Yes</u>	<u>DK</u>	
A1. Bizarre delusions	1	3	9	(184)
A2. Hallucinations	1	3	9	(185)
A3. Disorganized speech (frequent derailment or incoherence)	1	3	9	(186)
A4. Grossly disorganized or catatonic behavior	1	3	9	(187)
A5. Negative symptoms, i.e., affective flattening, alogia, or avolition	1	3	9	(188)

N.B. Only one Criterion A symptom is required if delusions are bizarre or hallucinations consist of a voice keeping up a running commentary on the person's behavior or thoughts, or two or more voices conversing with each other.

Criterion A:	1	3	9	(189)
If <u>any psychotic symptom</u> rated 2 or 3, (i.e., symptoms in A.1 through A7.3), code age at onset:		____		(190-191)
If <u>ALL ELEMENTS</u> of Criterion A1 through A5 all = 1, rate Schizophrenia, Schizoaffective Disorder and Brief Psychotic Disorder = 1, and skip to Obsessive-Compulsive Disorder.				

B. Social or occupational dysfunction 1 3 9 (192)

C. Continuous signs of disturbance for at least six months, which can include prodromal and residual periods. 1 3 9 (193)

PSYCHOSIS/SCHIZOPHRENIA (CONTINUED)

	<u>No</u>	<u>Yes</u>	<u>DK</u>	
D. Any affective syndrome has been short relative to active and residual phases	1	3	9	(194)
E. Not due to substance use or medical condition	1 due to organic factor	3 not due to organic factor	9	(195)
F. If there is history of Autistic Disorder or another Pervasive Development disorder, prominent delusions or hallucinations for at least one month. (Note: if no history of Autistic Disorder or another Pervasive Developmental Disorder, rate this criterion = 3)	1	3	9	(196)

Circle Schizophrenia = 3 if all of Criteria A through F above are coded 3.

SCHIZOPHRENIA.....1.....2.....3.....9 (197)

If Schizophrenia = 2 or 3: Age first sought treatment: _____ (198-199)

Age symptoms first caused distress or impairment: _____ (200-201)

Age first met criteria for schizophrenia: _____ (202-203)

SCHIZOAFFECTIVE DISORDER

	<u>No</u>	<u>Yes</u>	<u>DK</u>	
A. Either a Major Depressive, Manic or Mixed Episode(s) concurrent with symptoms that meet Criterion A of Schizophrenia	1	3	9	(204)
B. During an episode there have been delusions or hallucinations for at least two weeks, in the absence of prominent mood symptoms	1	3	9	(205)
C. Symptoms that meet criteria for mood episodes are present for a substantial portion of the total durations of the active and residual periods	1	3	9	(206)
D. Not due to substance use or medical disorder	1	3	9	(207)
E. Schizophrenia has been ruled out (i.e., Schizophrenia rating, pg. 15 = 1)	1	3	9	(208)

Circle Schizoaffective Disorder = 3 if all the above criteria A through E are coded 3.

SCHIZOAFFECTIVE DISORDER.....1.....2.....3.....9 (209)

If Schizoaffective Disorder = 2 or 3:

Age first sought treatment: _____ (210-211)

Age symptoms first caused distress or impaired functioning: _____

Circle Type: Depressive Type 1 3 9 (211a)

Bipolar Type 1 3 9 (211b)

BRIEF PSYCHOTIC DISORDER

	<u>No</u>	<u>Yes</u>	<u>DK</u>	
A. Presence of one or more of the following:	1	3	9	(212)
1. Delusions	1	3	9	(213)
2. Hallucinations	1	3	9	(214)
3. Disorganized Speech	1	3	9	(215)
4. Grossly Disorganized or Catatonic Behavior	1	3	9	(216)

CODE CRITERION A.

B. Duration from one day to one month, with eventual full return to premorbid level of functioning	1	3	9	(217)
C. No affective disorder, not due to medical condition or substance use	1	3	9	(218)

Circle Brief Psychotic Disorder = 3 if all the above criteria A through C are coded 3.

BRIEF PSYCHOTIC DISORDER.....1.....2.....3.....9 (219)

If Brief Psychotic Disorder = 2 or 3, age at onset: _____ (220-221)

OBSESSIVE COMPULSIVE DISORDER

A. Either obsessions (as defined by A.1, A.2, A.3 and A.4) or compulsions (as defined by A.5 and A.6):

	<u>No</u>	<u>Probable</u>	<u>Yes</u>	<u>D/K</u>
A.1 Recurrent and persistent thoughts, impulses, or images that are experienced, at some time during the disturbance, as intrusive and inappropriate and that cause marked anxiety or distress.	1	2	3	9
If <u>no</u> obsessions, skip to Compulsions (A-6) and rate Obsessions = 1				
A.2 The thoughts, impulses or images are not just excessive worries about real-life problems	1	2	3	9
A.3 Attempts to ignore or suppress thoughts, impulses, & images, or to neutralize them with other thought or action	1	2	3	9
A.4 The person recognizes as product of own mind	1	2	3	9

Elements of A.1

Aggressive Obsessions	1	2	3	9	(222)
Contamination Obsessions	1	2	3	9	(223)
Sexual Obsessions	1	2	3	9	(224)
Hoarding/Saving	1	2	3	9	(225)
Religious (Scrupulosity)	1	2	3	9	(226)
Need for Symmetry	1	2	3	9	(227)
Miscellaneous	1	2	3	9	(228)
Somatic	1	2	3	9	(229)

Presence of Obsessions.....1.....2.....3.....9 (230)

OCD (CONTINUED)

No Probable Yes D/K

Compulsions:

A.5 Repetitive Behaviors or mental acts that the person feels driven to perform in response to an obsession, or according to rules that must be applied rigidly

1 2 3 9

(If no Obsessions or Compulsions, Rate Obsessions and Compulsions = 1, and skip to Tic Disorders.)

A.6 The behaviors or mental acts are aimed at preventing or reducing distress or preventing some dreaded event or situation; behaviors or mental acts not connected in realistic way with what they are designed to prevent or neutralize or are clearly excessive

1 2 3 9

Cleaning/Washing Compulsions	1	2	3	9	(231)
Checking Compulsions	1	2	3	9	(232)
Repeating Rituals	1	2	3	9	(233)
Counting Compulsions	1	2	3	9	(234)
Ordering/Arranging Compulsions	1	2	3	9	(235)
Hoarding/Collecting Compulsions	1	2	3	9	(235a)
Miscellaneous Compulsions	1	2	3	9	(236)

Presence of Compulsions.....1.....2.....3.....9 (237)

OCD (CONTINUED)

	<u>No</u>	<u>Probable</u>	<u>Yes</u>	<u>DK</u>	
A.	Presence of Criterion A (either obsessions or compulsions)				(238)
	1	2	3	9	
B.	At some point during the course of the disorder, the person has recognized that the obsessions or compulsions are excessive or unreasonable (This does not apply to children).				(239)
	1	2	3	9	
C.	The obsessions or compulsions cause marked distress, are time consuming or significantly interfere with the person's normal routine, occupational (or academic) functioning, or usual social activities or relationships.				(240)
	1	2	3	9	
D.	If another Axis I disorder is present, the content of the obsessions or compulsions is not restricted to it				(241)
	1	2	3	9	
E.	The disturbance is not due to the direct physiological effects of a substance or a general medical condition.				(242)
	1	2	3	9	

Circle Obsessive/Compulsive Disorder=3
if all the above criteria A through E are coded 3

OBSESSIVE/COMPULSIVE DISORDER.....1.....2.....3.....9 (243)

If Obsessive/Compulsive Disorder = 2 or 3; or obsessions or compulsions = 2 or 3:

Age first experienced symptoms: _____ (256-257)

Age treatment first sought: _____ (258-259)

Age met diagnostic criteria: _____ (260-261)

TIC DISORDERS

		<u>No</u>	<u>Yes</u>	<u>DK</u>	
A.	Motor Tics:				
	1) Single	1	3	9	(263)
	2) Multiple	1	3	9	(264)
B.	Vocal Tics:				
	1) Single	1	3	9	(265)
	2) Multiple	1	3	9	(266)

(If no tics (ever), circle DSM-III-R and DSM-IV Tourette's Disorder = 1, DSM-III-R and DSM-IV Chronic Motor or Vocal Tics Disorder and DSM-IV Transient Tic disorder = 1 and SKIP to Hypochondriasis.)

C.	Both multiple motor <u>and</u> one or more vocal tics present.	1	3	9	(267)
D.	Tics occur many times per day, nearly every day or intermittently throughout a period of more than one year.	1	3	9	(268)
E.	As above <u>but</u> greater than 2 weeks and less than one year.	1	3	9	(269)
F.	Never a tic-free period of more than 3 consecutive months	1	3 never	9	(270)
G.	Marked distress or significant impairment	1	3	9	(271)
H.	Onset before 18	1	3	9	(272)
I.	Onset before 21	1	3	9	(273)
J.	Anatomic location, number, frequency and complexity of tics change over time	1	3	9	(274)
K.	<u>Not</u> due to medical or substance induction	1	3 not due to	9	(275)

Circle DSM-III-R Tourette's Disorder = 3, if Criteria C, D, I, and J = 3

DSM-III-R TOURETTE'S DISORDER.....1.....2.....3.....9 (276)

Circle DSM-IV Tourette's Disorder = 3, if Criteria C, D, F, G, and H = 3

DSM-IV TOURETTE'S DISORDER.....1.....2.....3.....9 (277)

Circle DSM-III-R Chronic Motor or Vocal Tic Disorder = 3, if Criteria A or B (not both), D, I, K = 3

DSM-III-R CHRONIC MOTOR OR VOCAL TIC DISORDER.....1.....2.....3.....9 (278)

TIC DISORDERS (CONTINUED)

Circle DSM-IV Chronic Motor or Vocal Tic Disorder = 3, if Criteria A or B (not both), D, F, G, H, K = 3

**DSM-IV CHRONIC MOTOR OR
VOCAL TIC DISORDER.....1.....2.....3.....9** (279)

Circle DSM-IV Transient Tic Disorder = 3, if Criteria A and/or B, E, G, H, K = 3

DSM-IV TRANSIENT TIC DISORDER.....1.....2.....3.....9 (280)

DSM-IV TIC DISORDER NOS 1.....2..... 3.....9 (280a)

For disorders characterized by tics that do not meet criteria for a specific Tic Disorder.
Ex: tics lasting less than 4 weeks, or tics with onset after 18 years old.

Age at Onset of any of above Tic Disorders _____ (281-282)

HYPOCHONDRIASIS

	<u>No</u>	<u>Yes</u>	<u>DK</u>	
A. Preoccupation with fear/belief that one has a serious disease	1	3	9	(283)
If Criteria A = 1, Circle Hypochondriasis = 1, and Skip to BDD				
B. Fear/belief persists despite medical reassurance	1	3	9	(284)
C. Belief in "A" not delusional <u>or</u> circumscribed (BDD)	1	3	9	(285)
D. Clinically significant distress/impairment	1	3	9	(286)
E. Duration ≥ 6 months	1	3	9	(287)
F. Not accounted for by GAD, OCD, Panic Disorder, MDD, Sep. Anxiety Disorder or another somatoform disorder	1	3	9	(288)

Circle Hypochondriasis = 3, if Criteria A through F = 3

HYPOCHONDRIASIS.....1.....2.....3.....9 (289)

Age of Onset if = 2 or 3: _____ (290-291)

BODY DYSMORPHIC DISORDER

	<u>No</u>	<u>Yes</u>	<u>DK</u>	
A. Preoccupation with an imagined defect in appearance. If slight, concern is markedly excessive.	1	3	9	(292)

If Criteria A = 1, Circle BDD = 1, and Skip to Trichotillomania

B. Clinically significant distress or impairment.	1	3	9	(293)
C. Not better accounted for by another mental disorder	1	3	9	(294)

Circle Body Dysmorphic Disorder = 3, if criteria A, B, and C = 3

BODY DYSMORPHIC DISORDER.....1.....2.....3.....9 (295)

Age of onset if = 2 or 3: _____ (296)

TRICHOTILLOMANIA

	<u>No</u>	<u>Yes</u>	<u>DK</u>	
A. Recurrent pulling out of one's hair resulting in noticeable hair loss.	1	3	9	(297)
If Criteria A = 1, Circle Trichotillomania = 1, and Skip to Nail Biting				
B. Tension immediately before pulling out the hair or resisting the behavior.	1	3	9	(298)
C. Pleasure, gratification, or relief when pulling out the hair.	1	3	9	(299)
D. Not accounted for by another mental disorder	1	3	9	(300)
E. Significant distress or impairment in social, occupational, or other important areas of functioning.	1	3	9	(301)

Circle Trichotillomania = 3, if Criterion A through E = 3

TRICHOTILLOMANIA.....1.....2.....3.....9 (302)

Age of onset, if 2 or 3: _____ (303-304)

NAIL BITING

	<u>No</u>	<u>Yes</u>	<u>DK</u>	
A. Recurrent nail biting resulting in noticeable evidence.	1	3	9	(305)
If Criteria A = 1, Circle Nail Biting = 1, and Skip to Skin Picking				
B. Tension immediately before nail biting or resisting the behavior.	1	3	9	(306)
C. Pleasure, gratification, or relief when biting nails.	1	3	9	(307)
D. Not accounted for by another mental disorder	1	3	9	(308)
E. Significant distress or impairment in social, occupational, or other important areas of functioning.	1	3	9	(309)

Circle Nail Biting = 3, if Criterion A through E = 3

NAIL BITING.....1.....2.....3.....9 (310)

Age of onset, if 2 or 3: (311-312)

SKIN PICKING

	<u>No</u>	<u>Yes</u>	<u>DK</u>	
A. Recurrent skin picking resulting in noticeable scarring.	1	3	9	(313)
If Criteria A = 1, Circle Skin Picking = 1, and Skip to Pathological Gambling				
B. Tension immediately before picking skin or resisting the behavior.	1	3	9	(314)
C. Pleasure, gratification, or relief when skin picking	1	3	9	(315)
D. Not accounted for by another mental disorder	1	3	9	(316)
E. Significant distress or impairment in social, occupational, or other important areas of functioning.	1	3	9	(317)

Circle Skin Picking = 3, if Criterion A through E = 3

SKIN PICKING.....1.....2.....3.....9 (318)

Age of onset, if 2 or 3: (319-320)

PATHOLOGICAL GAMBLING

A. Persistent and recurrent maladaptive gambling behavior as indicated by five (or more) of the following:

If doesn't Gamble, Circle Pathological Gambling = 1, and Skip to Pyromania

	<u>No</u>	<u>Yes</u>	<u>DK</u>	
(1) is preoccupied with gambling	1	3	9	(321)
(2) needs to gamble with increasing amounts of money	1	3	9	(322)
(3) repeated unsuccessful efforts to control gambling	1	3	9	(323)
(4) restless/irritable when attempting to cut down gambling	1	3	9	(324)
(5) gambles to escape problems or relieving a dysphoric mood	1	3	9	(325)
(6) gambling another day to get even	1	3	9	(326)
(7) lies to conceal gambling	1	3	9	(327)
(8) illegal acts to finance gambling	1	3	9	(328)
(9) lost relationship, job, etc. because of gambling	1	3	9	(329)
(10) relies on others to provide money because of gambling	1	3	9	(330)
CRITERION A: 5 of above met	1	3	9	(331)

B. The gambling behavior is not better accounted for by a Manic Episode. (332)

Circle Pathological Gambling = 3, if both Criterion A and B are coded 3.

PATHOLOGICAL GAMBLING.....1.....2.....3.....9 (333)

Age of onset, if 2 or 3: _____ (334-335)

PYROMANIA

	<u>No</u>	<u>Yes</u>	<u>DK</u>	
A. Deliberate and purposeful fire setting on more than one occasion.	1	3	9	(336)
If Criterion A = 1, Circle Pyromania = 1, and Skip to Kleptomania				
B. Tension or affective arousal before the act.	1	3	9	(337)
C. Fascination/curiosity/attraction/interest with fires	1	3	9	(338)
D. Pleasure, etc., when setting or witnessing fires	1	3	9	(339)
E. The fire setting not for monetary gain, criminal activity, anger or illness.	1	3	9	(340)
F. The fire setting is not better accounted for by Conduct Disorder, a Manic Episode, or Antisocial Personality Disorder.	1	3	9	(341)

Circle Pyromania = 3, if Criterion A through F = 3

PYROMANIA.....1.....2.....3.....9 (342)

Age of onset, if 2 or 3: _____ (343-344)

KLEPTOMANIA

	<u>No</u>	<u>Yes</u>	<u>DK</u>	
A. Recurrent failure to resist impulses to steal.	1	3	9	(345)
If Criterion A = 1, Circle Kleptomania = 1, and Skip to Panic Disorder				
B. Tension immediately before theft.	1	3	9	(346)
C. Pleasure, gratification, or relief at the time of committing the theft.	1	3	9	(347)
D. The stealing is not committed to express anger or due to psychosis.	1	3	9	(348)
E. Not accounted for by Conduct Disorder, a Manic Episode, or Antisocial Personality Disorder.	1	3	9	(349)

Circle Kleptomania = 3, if Criteria A through E = 3

KLEPTOMANIA.....1.....2.....3.....9 (350)

Age of onset, if 2 or 3: ____ ____ (351-352)

PANIC DISORDER

A. Enter number of symptoms in recurrent spontaneous panic attacks: _____ (99 = cannot be established) (353)

(If Criterion A = 0, circle Panic Disorder = 1 and Skip to Agoraphobia.)

	<u>No</u>	<u>Yes</u>	<u>DK</u>	
B. Symptoms started abruptly and peaked within 10 minutes	1	3	9	(354)
C. At least one of the attacks was followed by at least one month of persistent worry of another attack, worry about the implications of the attack or its consequences, or a significant change in behavior	1	3	9	(355)
D. Not due to medical condition or substance use	1	3	9	(356)
E. Attacks are not situation-specific (e.g., phobia)	1	3	9	(357)

Circle Panic Disorder = 3 if Criterion A >= 4 and Criteria B through E = 3.

PANIC DISORDER.....1.....2.....3.....9 (358)

If Panic Disorder = 2 or 3, age at onset: _____ (359-360)

SOCIAL PHOBIA

	<u>No</u>	<u>Yes</u>	<u>DK</u>	
A. Fear of social situation(s)	1	3	9	(371)
(If criterion A rated 1, circle Social Phobia = 1, and skip to Specific Phobia.)				
B. Exposure provokes anxiety	1	3	9	(372)
C. Avoidance or endured with intense anxiety	1	3	9	(373)
D. Interference with activities or marked distress about having fear	1	3	9	(374)
E. Recognizes fear as unreasonable or excessive	1	3	9	(375)
F. Unrelated to concerns about another medical or psychiatric disorder	1	3	9	(376)
G. Not due to medical condition or substance use	1	3	9	(377)

Circle Social Phobia = 3 if criteria A through G are met.

SOCIAL PHOBIA.....1.....2.....3.....9 (378)

If Social Phobia = 2 or 3, age at onset: _____ (379-380)

NOTE: Rate specific type of fear, even if subject does not meet criteria for phobia.

Code type:

Specific	1	3	9	(380a)
Generalized	1	3	9	(380b)

SPECIFIC PHOBIA

	<u>No</u>	<u>Yes</u>	<u>DK</u>	
A. Fear of circumscribed stimulus	1	3	9	(381)
(If criterion A rated 1, circle Specific Phobia = 1, and skip to General Anxiety Disorder.)				
B. Exposure provokes anxiety	1	3	9	(382)
C. Avoidance or endured with intense anxiety	1	3	9	(383)
D. Interference with activities or marked distress about having fear	1	3	9	(384)
E. Recognizes fear as unreasonable or excessive	1	3	9	(385)
F. Phobic stimulus is unrelated to content of another disorder, e.g., obsessions in OCD or the trauma of PTSD	1	3	9	(386)

Circle Specific Phobia = 3 if criteria A through F are met.

SPECIFIC PHOBIA.....1.....2.....3.....9 (387)

If Specific Phobia = 2 or 3, age at onset: _____ (388-389)

NOTE: Rate specific type of fear, even if subject does not meet criteria for phobia.

Code type:

Animal	1	3	9	(390)
Natural Environment	1	3	9	(391)
Blood / Injection / Injury	1	3	9	(392)
Situational	1	3	9	(393)
Other	1	3	9	(394)

GENERALIZED ANXIETY DISORDER

	<u>No</u>	<u>Yes</u>	<u>DK</u>	
A. Excessive anxiety and worry for at least a six-month period	1	3	9	(395)
(If criterion A= 1, circle Generalized Anxiety Disorder = 1 and go directly to Alcohol Dependence.)				
B. Subject finds it difficult to control the worry	1	3	9	(396)
C. The anxiety or worry is associated with at least three of the following six symptoms, with at least some symptoms present more days than not for at least six months	1	3	9	(397)

Elements of Criterion C:				
1. Restlessness or feeling keyed up or on edge	1	3	9	(398)
2. Being easily fatigued	1	3	9	(399)
3. Difficulty concentrating or mind going blank	1	3	9	(400)
4. Irritability	1	3	9	(401)
5. Muscle tension	1	3	9	(402)
6. Sleep disturbance	1	3	9	(403)
CODE CRITERION C				

D. The focus of the anxiety and worry is not confined to features of another Axis I disorder	1	3	9	(415)
E. The anxiety, worry, or physical symptoms cause clinically significant distress or impairment in social, occupational, or other important areas of functioning	1	3	9	(416)
F. Not due to medical condition or substance use, and not just occurring during episode of mood, psychotic, or pervasive developmental disorder	1	3	9	(417)

Circle Generalized Anxiety Disorder = 3 if Criteria A through F = 3.

GENERALIZED ANXIETY DISORDER.....1.....2.....3.....9 (418)

If Generalized Anxiety Disorder = 2 or 3, age at onset: _____ (419-420)

ALCOHOL DEPENDENCE

	<u>No</u>	<u>Yes</u>	<u>DK</u>	
Evidence of excessive alcohol use or alcohol-related problems	1	3	9	(421)

If no evidence circle Alcohol Dependence and Alcohol Abuse = 1 and skip to Substance Dependence

A. At least three of the following occurred at any time in the same 12-month period.	1	3	9	(422)
--	---	---	---	-------

Elements of Criterion A:				
1. Alcohol taken in larger amount or over longer period than the subject intended (1)	1	3	9	(423)
2. Persistent desire or one or more unsuccessful efforts to cut down or control substance use (2)	1	3	9	(424)
3. Marked tolerance (3a or b)	1	3	9	(425)
4. Either or both of the following: (4)	1	3	9	(426)
a. Withdrawal symptoms	1	3	9	(427)
b. Use to avoid withdrawal	1	3	9	(428)
5. Use while knowing physical and psychological sequelae (5)	1	3	9	(429)
6. Social, occupational, and recreational activities affected (6 and impairment)	1	3	9	(430)
7. Excessive time spent getting, taking, or recovering from effects (7)	1	3	9	(431)
CODE CRITERION A				

Circle Alcohol Dependence = 3 if Criterion A is = 3.

ALCOHOL DEPENDENCE.....1.....2.....3.....9 (432)

If Alcohol Dependence = 2 or 3, age at onset: _____ (433-434)

If Alcohol Dependence is circled "3", go directly to Substance Dependence ratings. Otherwise, continue with Alcohol Abuse ratings.

ALCOHOL ABUSE

	<u>No</u>	<u>Yes</u>	<u>DK</u>	
A. Continued use despite social and interpersonal consequences (5)	1	3	9	(435)
B. Hazardous use (9)	1	3	9	(436)
C. Recurrent substance-related legal problems (8)	1	3	9	(437)
D. Recurrent use resulting in failure to fulfill major role obligations (6)	1	3	9	(438)

Circle Alcohol Abuse = 3 if one or more of the criteria A through D led to clinically significant impairment or distress, within a 12-month period.

ALCOHOL ABUSE.....1.....2.....3.....9 (439)

Alcohol Abuse = 2 or 3, age at onset: _____ (440-441)

SUBSTANCE DEPENDENCE

	<u>No</u>	<u>Yes</u>	<u>DK</u>	
Evidence of excessive substance use <u>or</u> substance-related problems	1	3	9	(442)

If no evidence, circle Substance Dependence and Abuse = 1 and Skip to Anorexia Nervosa

If yes, circle the substance causing the most problems:

1. **Cannabis:** marijuana, hashish, THC, or other
2. **Cocaine:** intranasal, IV, freebase, crack, "speedball," unspecified or other
3. **Stimulants:** amphetamine, "speed," crystal meth, dexadrine, Ritalin, "ice", or other
4. **Sedatives-hypnotics-anxiolytics:** Quaalude, Seconal, Valium, Xanax, Librium, barbiturates, Miltown, Ativan, Dalmane, Halcion, Restoril, or other
5. **PCP ("Angel Dust")**
6. **Hallucinogens:** LSD, mescaline, peyote, psilocybin, STP, mushrooms, Extasy, MDMA, or other
7. **Opioids:** heroin, morphine, opium, Methadone, Darvon, codeine, Percodan, Demerol, Dilaudid, unspecified or other
8. **Inhalants:** "glue", paint, inhalants, nitrous oxide ("laughing gas"), amyl or butyl nitrate ("poppers")
9. **Polydrug**

SUBSTANCE DEPENDENCE (CONTINUED)

	<u>No</u>	<u>Yes</u>	<u>DK</u>	
A. At least three of the following occurred at any time in the same 12-month period.	1	3	9	(452)
Elements of Criterion A:				
1. Substance taken in larger amount or over longer period than the subject intended (1)	1	3	9	(453)
2. Persistent desire or one or more unsuccessful efforts to cut down or control substance use (2)	1	3	9	(454)
3. Marked tolerance (3)	1	3	9	(455)
4. Either of the following:	1	3	9	(456)
a. Withdrawal symptoms (4a)	1	3	9	(457)
b. Use to avoid withdrawal (4b)	1	3	9	(458)
5. Use while knowing physical, social, and psychological sequelae (5)	1	3	9	(459)
6. Social, occupational, and recreational activities affected (6 and impairment)	1	3	9	(460)
7. Excessive time spent getting, taking, or recovering from effects (7)	1	3	9	(461)
CODE CRITERION A				

Circle Substance Dependence = 3 if criterion A = 4 for either or both of two substances.

SUBSTANCE DEPENDENCE.....1.....2.....3.....9 (472)

If Substance Dependence = 2 or 3, age at onset: _____ (473-474)

If Substance Dependence is circled "3", go directly to Anorexia Nervosa. Otherwise, continue with Substance Abuse ratings.

SUBSTANCE ABUSE

Rate for up to 2 substances that did not receive a Substance Dependence rating of “3”.

Circle Substance Code Number: 1 2 3 4 5 6 7 8 9

	<u>No</u>	<u>Yes</u>	<u>DK</u>	
A. Use while knowing social and interpersonal consequences (10)	1	3	9	(475)
B. Hazardous use (8)	1	3	9	(476)
C. Recurrent substance-related legal problems (9)	1	3	9	(477)
D. Recurrent use resulting in failure to fulfill major role obligations (6)	1	3	9	(478)

Circle Substance Abuse = 3 if one or more of Criteria A through D are met for either or both of two substances.

SUBSTANCE ABUSE.....1.....2.....3.....9 (483)

If Substance Abuse = 2 or 3, age at onset: _____ (484-485)
(for earliest substance)

ANOREXIA NERVOSA

	<u>No</u>	<u>Yes</u>	<u>DK</u>	
A. Refusal to maintain body weight at = 85% of expected	1	3	9	(486)
If Criterion A = 1, circle Anorexia Nervosa = 1, and skip to Bulimia Nervosa.				
B. Intense fear of gaining weight or becoming fat	1	3	9	(487)
C. Disturbance in body perception	1	3	9	(488)
D. <u>Females</u> : absence of at least three consecutive menstrual cycles when otherwise expected to occur	1	3	9	(489)

**Circle Anorexia Nervosa = 3 if: Females: criteria A through D are met.
Males: criteria A through C are met**

ANOREXIA NERVOSA.....1.....2.....3.....9 (490)

If Anorexia Nervosa = 2 or 3, age at onset: _____ (491-492)

Code Type:

Restricting Type	1	3	9	(493)
Binge-eating / Purging type	1	3	9	(494)

BULIMIA NERVOSA

	<u>No</u>	<u>Yes</u>	<u>DK</u>	
A. Recurrent episodes of binge eating (rapid consumption of large amounts of food in a discrete period of time)	1	3	9	(495)
If Criterion A = 1, circle Bulimia Nervosa = 1, and skip to Dementia.				
B. Feeling of lack of control over eating behavior during the eating binges	1	3	9	(496)
C. Subject regularly engages in either self-induced vomiting, use of laxatives or diuretics, strict dieting, or fasting, or vigorous exercise in order to prevent weight gain	1	3	9	(497)
D. A minimum average of two binge eating episodes a week for at least three months	1	3	9	(498)
E. Persistent overconcern with body shape and weight	1	3	9	(499)
F. Symptoms do not occur exclusively during episodes of Anorexia Nervosa	1	3	9	(500)

Circle Bulimia Nervosa = 3 if criteria A through F are met.

BULIMIA NERVOSA.....1.....2.....3.....9 (501)

If Bulimia Nervosa = 2 or 3, age at onset: _____ (502-503)

Code Type:

Purging Type	1	3	9	(504)
Nonpurging Type	1	3	9	(505)

PTSD

No

Yes

DK

A. The subject has been exposed to a traumatic event, which involved threat to physical integrity, etc., AND the response involved intense fear, etc.

1

3

9

(534)

(If Criterion A = 1, circle PTSD = 1 and go to next disorder)

B. At least one of the following (reexperiencing):

1

3

9

(535)

1. Recurrent and intrusive recollections

1

3

9

(536)

2. Recurrent distressing dreams

1

3

9

(537)

3. Acting or feeling like event recurring

1

3

9

(538)

4. Intense distress with cues

1

3

9

(539)

5. Physiological reactivity with cues

1

3

9

(540)

CODE CRITERION B

C. At least three of the following (avoidance):

1

3

9

(541)

1. Efforts to avoid thoughts/feelings/conversations

1

3

9

(542)

2. Efforts to avoid activities/places/people

1

3

9

(543)

3. Inability to recall important aspect of trauma

1

3

9

(544)

4. Diminished interest or participation in activities

1

3

9

(545)

5. Feeling of detachment or estrangement from others

1

3

9

(546)

6. Restricted range of affect (e.g., no loving feelings)

1

3

9

(547)

7. Sense of foreshortened future

1

3

9

(548)

CODE CRITERION C

PTSD (CONTINUED)

D. At least two of the following (increased arousal): 1 3 9 (549)

1. Difficulty falling or staying asleep	1	3	9	(550)
2. Irritability or outbursts of anger	1	3	9	(551)
3. Difficulty concentrating	1	3	9	(552)
4. Hypervigilance	1	3	9	(553)
5. Exaggerated startle response	1	3	9	(554)
CODE CRITERION D				

E. Duration of symptoms is more than one month. 1 3 9 (555)

F. The disturbance causes clinically significant distress or impairment in social, occupational, or other important areas of functioning. 1 3 9 (556)

Circle PTSD = 3 if all the Criteria A through F = 3.

PTSD.....1.....2.....3.....9 (557)

If PTSD 2 or 3, code age at onset: _____ (558-559)

OTHER AXIS I DIAGNOSES:

No

Prob.

Definite

DK

Disorder: _____ 1 2 3 9 (543)

DSM-IV Code: ____ . ____

Disorder: _____ 1 2 3 9 (544)

DSM-IV Code: ____ . ____

Disorder: _____ 1 2 3 9 (544a)

DSM-IV Code: ____ . ____

Disorder: _____ 1 2 3 9 (544b)

DSM-IV Code: ____ . ____

Disorder: _____ 1 2 3 9 (544c)

DSM-IV Code: ____ . ____

AT THAT TIME, WHEN YOU WERE (AGE -AT WHICH SYMPTOMS WERE THE WORST)...

"I am now going to ask several questions about your obsessive thoughts."
 [Make specific reference to the patient's target obsessions.]

**1. How much of your time was occupied by obsessive thoughts?
 How frequently did they occur?**

____(665)

- 0=No interference
- 1=Mild, occasional intrusion (less than 1 hr/day)
- 2=Moderate, frequent intrusion (1 to 3 hrs/day)
- 3=Severe, very frequent intrusion (greater than 3 to 8 hrs/day)
- 4=Extreme, nearly constant intrusion (greater than 8 hrs/day)

2. How much did your obsessions interfere with your social, work or major role of functioning? Is there anything you didn't do because of them? [If you were not working, how much would your performance be affected if you were working?]

____(666)

- 0=No interference
- 1=Mild, slight interference with social or occupational activities, but overall performance not impaired
- 2=Moderate, definite interference with social or occupational performance, but still manageable
- 3=Severe, caused substantial impairment in social or occupational performance
- 4=Extreme, incapacitating

***3. How much distress did your obsessive thoughts cause you?**

- 0=No distress
- 1=Mild, infrequent and not too disturbing distress
- 2=Moderate, frequent, and disturbing, but still manageable distress
- 3=Severe, very frequent, and very disturbing distress
- 4=Extreme, near constant, and disabling distress

OBSESSION-FREE INTERVAL

4. **On the average, what is the longest number of consecutive waking hours per day that you are completely free of obsessive thoughts? [If necessary, ask:] What is the longest block of time in which obsessive thoughts are absent?** _____ (668)

0 = No symptoms.

1 = Long symptom-free interval, more than 8 consecutive hours/day symptom-free.

2 = Moderately long symptom-free interval, more than 3 and up to 8 consecutive hours/day symptom-free.

3 = Short symptom-free interval, from 1 to 3 consecutive hours/day symptom-free.

4 = Very short symptom-free interval, less than 1 consecutive hour/day symptom-free; freedom from obsessions measured in minutes.

5 = No symptom-free interval, constant obsessions without even momentary relief.

- *5. **How much of an effort did you make to resist your obsessions? How often did you try to disregard or turn your attention away from obsessions as they entered your mind? [Rate only the effort you make to resist, not your success or failure in actually controlling the obsessions.]** _____ (668)

0=Always tried to resist, or symptoms so minimal I didn't need to actively resist

1=Tried to resist most of the time

2=Made some effort to resist

3=Yielded to all obsessions without attempting to control them, but with some reluctance

4=Completely and willingly yielded to all obsessions

- *6. **How much control did you have over your obsessive thoughts? How successful were you in stopping or diverting your obsessive thinking?** _____ (669)

0=Complete control

1=Much control, usually was able to stop or divert obsessions with some concentration and effort

2=Moderate control, sometimes was able to stop or divert obsessions

3=Little control, rarely was successful in stopping obsessions, could only divert attention with difficulty

4=No control, obsessions were experienced as completely involuntary, rarely was able to even momentarily divert thinking

7. **How much time did you spend performing compulsive behaviors (rituals)? How frequently were they performed? How much longer than most people did it take to complete routine activities because of your rituals? [Many rituals are observable behaviors (e.g. hand washing), but others are not observable (e.g. mental checking or silent safety statements such as, "God is good. He will not let me harm my child.").]** _____ (670)

0=None

1=Mild, less than 1 hr/day performing rituals or occasional performance of rituals

2=Moderate, 1 to 3 hrs/day performing rituals, or frequent performance of rituals

3=Severe, between 3 to 8 hrs/day performing rituals or very frequent performance of rituals

4=Extreme, more than 8 hrs/day performing rituals, or near constant performance of rituals

8. How much did your rituals interfere with your social, work or major role of functioning? Is there anything you didn't do because of your rituals? [If you were not working then, how much would your performance be affected if you were?]

___(671)

0=None

1=Mild, slight interference with social or occupational performance, but overall performance was not impaired

2=Moderate, definite interference with social or occupational performance, but was still manageable

3=Severe, caused substantial impairment in social or occupational performance

4=Extreme, incapacitating

***9. How would you have felt if you were prevented from performing your rituals? How anxious would you have become? If having an outsider interrupt your rituals is actually agreeable and helpful, then answer this question with regard to how anxious you became until you were satisfied your rituals were properly completed.**

___ (672)

0=No distress

1=Mild, only slightly anxious if rituals were prevented, or only slight anxiety during performance of rituals

2=Moderate, anxiety would mount but would remain manageable if rituals were prevented, or that anxiety increased to moderate levels during performance of rituals

3=Severe, prominent and very disturbing increase in anxiety if rituals were interrupted, or prominent and very disturbing increase in anxiety during performance of rituals

4=Extreme, incapacitating anxiety from any intervention aimed at modifying rituals, or incapacitating anxiety developed during performance of rituals

10. How much effort did you make to resist your rituals? [Rate only effort to resist rituals, not your success or failure.]

___ (673)

0=Always resisted compulsions/rituals, or symptoms so minimal I didn't need to actively resist

1=Tried to resist rituals most of the time

2=Made some effort to resist compulsions/rituals

3=Yielded to all compulsions/rituals without attempting to control them, but with some reluctance

4=Completely and willingly yielded to all compulsions/rituals

COMPULSION-FREE INTERVAL

11. On the average, what is the longest number of consecutive waking hours per day that you are completely free of compulsive behaviors or avoiding situations or cues that might trigger your obsessions? [If necessary, ask:] What is the longest block of time in which compulsive or avoidant behaviors are absent?

___ (674)

0 = No symptoms.

1 = Long symptom-free interval, more than 8 consecutive hours/day symptom-free.

2 = Moderately long symptom-free interval, more than 3 and up to 8 consecutive hours/day symptom-free.

3 = Short symptom-free interval, from 1 to 3 consecutive hours/day symptom-free.

4 = Very short symptom-free interval, less than 1 consecutive hour/day symptom-free, freedom from compulsions **and/or avoidance** measured in minutes.

5 = No symptom-free interval, constant compulsions without even momentary relief.

***12. How strong was the drive to perform your rituals? How much control did you have over your rituals?**

____(674a)

0=Complete control

1=Much control, experienced pressure to perform rituals, but usually was able to exercise voluntary control over it

2=Moderate control, strong pressure to perform rituals, could control it only with difficulty

3=Little control, very strong drive to control rituals. Must have been carried out to completion, could only delay with difficulty

4=No control, drive to ritualize was completely involuntary and overpowering, was rarely able to even momentarily delay activity

13. Did you think your concerns or behaviors were reasonable? What did you think would happen if you did not perform the compulsion(s)? Were you convinced something would really happen? [Rate patient's insight into the senselessness or excessiveness of his obsession(s) based on beliefs expressed at the time of the interview.]

____(675)

0=Excellent insight, fully rational

1=Good insight. Readily acknowledged absurdity or excessiveness of thoughts or behaviors but did not seem to be completely convinced that there was something besides anxiety to be concerned about (i.e., has lingering thoughts).

2=Fair insight. Reluctantly admitted that thoughts or behaviors seemed unreasonable or excessive, but

3=Poor insight. Maintained that thoughts and behaviors were not unreasonable or excessive, but acknowledged validity of contrary evidence (i.e., overvalued ideas present).

4=Lacked insight, delusional. Definitely was convinced that concerns and behavior were reasonable, was unresponsive to contrary evidence.

14. Did you avoid doing anything, going any place, or being with anyone because of your obsessional thoughts or out of concern you would perform compulsions? [If yes, then ask:] How much did you avoid? [Rate degree to which patient deliberately tried to avoid things. Sometimes compulsions are designed to "avoid" contact with something that the patient fears. For example, clothes washing rituals would be designated as compulsions, not as avoidant behavior. If the patient stopped doing the laundry then this would constitute avoidance.]

____ (676)

0=No deliberate avoidance

1=Mild, minimal avoidance

2=Moderate, some avoidance; clearly present

3=Severe, much avoidance; avoidance prominent

4=Extreme, very extensive avoidance; patient did almost everything he/she could to avoid triggering symptoms

15. Did you have trouble making decisions about little things that other people might not think twice about (e.g., which clothes to put on in the morning; which brand of cereal to buy)? [Exclude difficulty making decisions which reflect ruminative thinking. Ambivalence concerning rationally-based difficult choices should also be excluded.]

____ (677)

0=None

1=Mild, some trouble making decisions about minor things

2=Moderate, freely reports significant trouble making decisions that others would not think twice about

3=Severe, continual weighing of pros and cons about nonessentials

4=Extreme, unable to make any decisions. Disabling.

- 16. Did you feel very responsible for the consequences of your actions? Did you blame yourself for the outcome of events not completely in your control? [Distinguish from normal feelings of responsibility, feeling of worthlessness, and pathological guilt. A guilt-ridden person experiences himself or his actions as bad or evil.]** _____(678)

0=None

1=Mild, only mentioned on questioning, slight sense of over-responsibility

2=Moderate, ideas stated spontaneously, clearly present; patient experienced significant sense of over-responsibility for events outside his/her reasonable control

3=Severe, ideas prominent and pervasive; deeply concerned he/she is responsible for events clearly outside his control. Self-blaming farfetched and nearly irrational

4=Extreme, delusional sense of responsibility (e.g., if an earthquake occurs 3,000 miles away patient blames herself because she didn't perform her compulsions)

- 17. Did you have difficulty starting or finishing tasks? Did many routine activities take longer than they should? [Distinguish from psychomotor retardation secondary to depression. Rate increased time spent performing routine activities even when specific obsessions cannot be identified.]** _____(679)

0=None

1=Mild, occasional delay in starting and finishing

2=Moderate, frequent prolongation of routine activities but tasks usually completed. Frequently late.

3=Severe, pervasive and marked difficulty initiating and completing routine tasks. Usually late.

4=Extreme, unable to start or complete routine tasks without full assistance.

- 18. After you complete an activity do you doubt whether you performed it correctly? Do you doubt whether you did it at all? When carrying out routine activities did you feel you didn't trust your senses (i.e., what you see, hear or touch)?** _____(680)

0=None

1=Mild, only mentioned on questioning, slight pathological doubt. Examples given may be within normal range

2=Moderate, ideas stated spontaneously, clearly present and apparent in some of patient's behaviors; patient bothered by significant pathological doubt. Some effect on performance but still manageable

3=Severe, uncertainty about perceptions or memory prominent; pathological doubt frequently affects performance

4=Extreme, uncertainty about perceptions constantly present; pathological doubt substantially affects almost all activities. Incapacitating (e.g., patient states "my mind doesn't trust what my eyes see").

- 19. Interviewer's judgement of the overall severity of the patient's illness. Rated from 0 (no illness) to 6 (most severe patient seen). [Consider the degree of distress reported by the patient, the symptoms observed, and the functional impairment reported. Your judgement is required both in averaging this data as well as weighing the reliability or accuracy of the data obtained. This judgement is based on information obtained during the interview.]** _____(681)

0=No illness

1=Illness slight, doubtful, transient; no functional impairment

2=Mild symptoms, little functional impairment

3=Moderate symptoms, functions with effort

4=Moderate - Severe symptoms, limited functioning

5=Severe symptoms, functions mainly with assistance

6=Extremely severe symptoms, completely nonfunctional